

WEIGHTED LEAST ABSOLUTE DEVIATIONS ESTIMATION FOR ARMA MODELS WITH INFINITE VARIANCE

JIAZHU PAN HUI WANG QIWEI YAO

Peking University and London School of Economics

For autoregressive and moving-average (ARMA) models with infinite variance innovations, quasi-likelihood based estimators (such as Whittle's estimators) suffer from complex asymptotic distributions depending on unknown tail indices. This makes the statistical inference for such models difficult. In contrast, the least absolute deviations estimators (LAD) are more appealing in dealing with heavy tailed processes. In this paper, we propose a weighted least absolute deviations estimator (WLAD) for ARMA models. We show that the proposed WLAD is asymptotically normal, unbiased and with the standard root- n convergence rate even when the variance of innovations is infinity. This paves the way for the statistical inference based on asymptotic normality for heavy-tailed ARMA processes. For relatively small samples numerical results illustrate that the WLAD with appropriate weight is more accurate than the Whittle estimator, the quasi-maximum likelihood estimator (QMLE) and the Gauss-Newton estimator when the innovation variance is infinite, and that the efficiency-loss due to the use of weights in estimation is not substantial.

The authors thank the two referees for their valuable suggestions. The work was partially supported by an EPSRC research grant (UK), and the Natural Science Foundation of China (grant 10471005). Address correspondence to Jiazhu Pan, School of Mathematical Sciences, Peking University, Beijing 100871, China; e-mail: jzpan@math.pku.edu.cn.

1 INTRODUCTION

Let $\{y_t\}$ be a stationary ARMA time series generated by the equation

$$y_t = \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p} + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q}, \quad (1.1)$$

where the innovation process $\{\varepsilon_t\}$ is a sequence of independent and identically distributed (iid) random variables, and $\beta = (\phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q)'$ is an unknown parameter vector. When $E\varepsilon_t^2 < \infty$, it is well known that various estimators such as MLE, Whittle's estimators and least squares estimators (LSE) for β are all asymptotically normal and unbiased (Brockwell and Davis 1991). When $E\varepsilon_t^2 = \infty$, model (1.1) is called the infinite variance ARMA (IVARMA) model which defines a heavy-tailed process $\{y_t\}$. The IVARMA models are pertinent in modelling heavy-tailed time series data often encountered in, for example, economics and finance (Koedijk et al. 1990, and Jansen and de Vries 1991). For further references on statistical modelling for heavy-tailed phenomena, we refer to Resnick (1997) and Adler et al. (1997).

Statistical inference for IVARMA models has not been well explored yet. Most available results concern with infinite variance AR (IVAR) models (i.e. $q = 0$ in (1.1)). Gross and Steiger (1979) and An and Chen (1982) obtained the strong consistency and the convergence rates for LADE for IVAR models. Davis and Resnick (1985, 1986) derived the limiting distributions of the LSE for IVAR models. A more comprehensive asymptotic theory of M-estimators for AR models was derived by Davis et al. (1992). For IVARMA models, the asymptotic properties for several estimators have been derived when the innovation distribution is in the domain of attraction of a stable law distribution with index between 0 and 2. For example, both the Whittle estimator proposed by Mikosch et al. (1995) and the Gauss-Newton estimator proposed by Davis (1996) converge in distribution to some functions of a sequence of stable random variables. Furthermore, Davis (1996) proved that the M-estimator converges in distribution to the minimizer of a stochastic process. However, all the limiting distributions above are complicated and depend intimately on the unknown tail indices of the underlying processes. This makes it difficult to develop asymptotic approximations for the purpose of statistical inference. This paper provides a remedy for this problem.

The difficulty of the asymptotic theory for LADE for IVARMA processes may at least par-

tially attribute to the fact that the residual of a linear prediction for y_t based on its lagged values depends on β *nonlinearly*, while such a dependence is completely linear for pure AR processes. Note that this linearity implies that the objective function for LAD estimation is convex, and therefore the asymptotic normality of LADE may be readily derived from the convex lemma (Hjort and Pollard 1993). One way to deal with a non-convex objective function is to adopt a local linearization around the true value of the parameter, which enables one to establish asymptotic properties of a local estimator defined as a local minimizer around the true value of the parameter. This is the line taken by Davis and Dunsmuir (1997) which dealt with LADE for ARMA models with $E\varepsilon_t^2 < \infty$. On the other hand, Ling (2005) proposed a WLADE for IVAR models. The key idea of the WLADE is to weigh down the observations which are excessively large, either positively or negatively. Ling (2005) showed that the WLADE is asymptotically normal. The idea of weighing down the large observations has also been used in estimation for ARCH models with heavy tailed innovations by Horvath and Liese (2004).

In this paper, we deal with the WLADE for IVARMA models. By adopting the idea of local linearization mentioned above, we show that a local WLADE is asymptotic normal and unbiased under the condition that $E|\varepsilon_t|^\delta < +\infty$ for some $\delta > 0$ and the density function of ε_t and its derivative are bounded. This facilitates the statistical inference for IVARMA models (even when $E|\varepsilon_t| = \infty$) in a conventional fashion. For example, a Wald test for a linear hypothesis can be constructed; see Section 2 below. For relatively small samples a simulation study indicates that the proposed WLADE is more accurate than the Whittle estimator, the quasi-maximum likelihood estimator (QMLE) and the Gauss-Newton estimator when $Var(\varepsilon_t) = \infty$. Furthermore, the efficiency-loss of the WLADE with respect to the (unweighted) LADE is not significant with appropriately selected weights. Since the WLADE converges at a slower rate than Whittle estimator and Gauss-Newton estimator, we also studied the large sample properties of the WLADE numerically.

Although we only deal with IVARMA models in this paper, the basic idea of combining a weighted objective function with local linearization of the residuals may apply to other infinite variance time series models, such as the infinite variance ARIMA and the integrated GARCH (IGARCH) which are popular in financial econometrics. Another open problem is to develop

appropriate methods for choosing weight functions; see Remark 3 below.

The rest of paper is organized as follows. The WLADE and the associated asymptotic properties are presented in section 2. In addition to the asymptotic normality of the local WLADE, we also show that a (global) estimator sharing the same asymptotic property could be obtained by minimizing a convex objective function if available an initial estimator within root- n distance from the true value; see Theorem 2 below. Section 3 gives all theoretical proofs of the results in section 2. Section 4 reports some numerical results from a simulation study.

2 WLADE AND ITS ASYMPTOTIC PROPERTIES

2.1 Weighted least absolute deviations estimators

Denote $\Theta \subset R^{p+q}$ the parameter space which contains the true value $\beta_0 = (\phi_1^0, \dots, \phi_p^0, \theta_1^0, \dots, \theta_q^0)'$ of the parameter β as an inner point. For $\beta = (\phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q)'$, put

$$\varepsilon_t(\beta) = \begin{cases} y_t - \phi_1 y_{t-1} - \dots - \phi_p y_{t-p} - \theta_1 \varepsilon_{t-1}(\beta) - \dots - \theta_q \varepsilon_{t-q}(\beta), & \text{if } t > 0, \\ 0, & \text{otherwise,} \end{cases} \quad (2.1)$$

where $y_t \equiv 0$ for all $t \leq 0$. Note that $\varepsilon_t \neq \varepsilon_t(\beta_0)$ due to this truncation.

We define the objective function as

$$W_n(\beta) = \sum_{t=u+1}^n \tilde{w}_t |\varepsilon_t(\beta)|, \quad (2.2)$$

and the weighted least absolute deviation estimator (WLADE) as

$$\hat{\beta} = \arg \min_{\beta} W_n(\beta) \quad (2.3)$$

where $u = u(n)$ is a positive integer, and the weight function, depending on a constant $\alpha > 2$, is defined as

$$\tilde{w}_t = \left(1 + \sum_{k=1}^{t-1} k^{-\alpha} |y_{t-k}|\right)^{-2}. \quad (2.4)$$

2.2 Asymptotic normality of WLADe

To state the asymptotic normality of $\hat{\beta}$, we introduce some notation first. Let

$$v = (v_1, \dots, v_{p+q})' = \sqrt{n}(\beta - \beta_0).$$

It is easy to see that $\hat{\beta} = \beta_0 + \hat{v}/\sqrt{n}$, where \hat{v} is a minimizer of

$$T_n(v) = \sum_{t=u+1}^n \tilde{w}_t(|\varepsilon_t(\beta_0 + n^{-1/2}v)| - |\varepsilon_t(\beta_0)|).$$

Denote $A_t(\beta) = (A_{t,1}(\beta), \dots, A_{t,p+q}(\beta))'$, where $A_{t,i}(\beta) = -\partial\varepsilon_t(\beta)/\partial\beta_i$. By (8.11.9) of Brockwell and Davis (1991), it holds for $t \geq \max(p, q)$ that

$$\begin{cases} \theta(B)A_{t,i}(\beta) = y_{t-i}, & i = 1, \dots, p \\ \theta(B)A_{t,i+p}(\beta) = \varepsilon_{t-i}(\beta), & i = 1, \dots, q, \end{cases} \quad (2.5)$$

where B is the backshift operator.

For $t = 0, \pm 1, \pm 2, \dots$, define

$$U_t - \sum_{i=1}^p \phi_i^0 U_{t-i} = \varepsilon_t, \quad V_t + \sum_{j=1}^q \theta_j^0 V_{t-j} = \varepsilon_t. \quad (2.6)$$

Put $Q_t = (U_{t-1}, \dots, U_{t-p}, V_{t-1}, \dots, V_{t-q})'$, $w_t \equiv (1 + \sum_{k=1}^{\infty} k^{-\alpha} |y_{t-k}|)^{-2}$, and

$$\Sigma = E(w_t Q_t Q_t'), \quad \Omega = E(w_t^2 Q_t Q_t'), \quad (2.7)$$

We denote by $\|v\|$ the Euclidean norm for a vector v .

Some regularity conditions are now in order.

A1 For $\beta \in \Theta$, the polynomials $\theta(z) = 1 + \theta_1 z + \dots + \theta_q z^q$ and $\phi(z) = 1 - \phi_1 z - \dots - \phi_p z^p$ have no common zeroes, and all roots of $\phi(z)$ and $\theta(z)$ are outside the unit circle.

A2 Innovation ε_t has zero median and a differentiable density function $f(x)$ satisfying the conditions $f(0) > 0$, $\sup_{x \in R} |f(x)| < B_1 < \infty$ and $\sup_{x \in R} |f'(x)| < B_2 < \infty$. Furthermore, $E|\varepsilon_t|^\delta < +\infty$ for some $\delta > 0$, and $\alpha > \max\{2, \frac{2}{\delta}\}$.

A3 As $n \rightarrow \infty$, $u \rightarrow \infty$ and $u/n \rightarrow 0$.

The following proposition indicates that model (1.1) has a unique strictly stationary and ergodic solution under condition A1-A2.

Proposition 1. *Suppose that condition A1 hold and $E|\varepsilon_t|^\delta < +\infty$ for some $\delta > 0$. Then model (1.1) defines a unique strictly stationary and ergodic process $\{y_t\}$.*

Proof. Condition A1 implies that

$$\phi^{-1}(z)\theta(z) = \sum_{j=0}^{\infty} \psi_j z^j,$$

where $|\psi_j| \leq Cr^j$ for some constants $C > 0$ and $0 < r < 1$. Let $\tilde{\delta} = \min\{\delta, 1\}$. Then

$$\sum_{j=0}^{\infty} |\psi_j|^{\tilde{\delta}} E|\varepsilon_{t-j}|^{\tilde{\delta}} < \infty.$$

The same argument for Proposition 13.3.2 of Brockwell and Davis (1991) yields the result. \square

Remark 1. Condition A3 eliminates asymptotically the bias in the estimation due to the lack of observations y_t for $t \leq 0$.

Remark 2. Condition A2 does not rule out the possibility that $E|\varepsilon_t| = \infty$. The purpose of introducing weights \tilde{w}_t is to weigh down excessively large observations which reflect the heavy-tailed innovation distribution. Therefore the asymptotic covariance matrix of the normalized WLAD, depending on Σ and Ω given in (2.7), is a well-defined (finite) matrix. Note that $\tilde{w}_t \in (0, 1]$. Conditions A1 and A2 imply that for $\tilde{\delta} = \min\{\delta, 1\}$,

$$E\left(\sum_{k=1}^{\infty} k^{-\alpha} |y_{t-k}|\right)^{\tilde{\delta}} \leq \sum_{k=1}^{\infty} k^{-\alpha\tilde{\delta}} E|y_{t-k}|^{\tilde{\delta}} < +\infty.$$

Hence $\sum_{k=1}^{\infty} k^{-\alpha} |y_{t-k}| < \infty$ with probability one, which ensures that w_t is well defined. Note that w_t is stationary and ergodic under Condition A1, and it is asymptotically equivalent to \tilde{w}_t for $t > u$ (see A3).

We are now ready to state our main results.

Theorem 1. *Let conditions A1 – A3 hold. For any given positive random variable M with $P(0 < M < \infty) = 1$, there exists a local minimizer $\hat{\beta}$ of $W_n(\beta)$ which lies in the random region $\{\beta : \|\beta - \beta_0 - \xi/\sqrt{n}\| \leq M/\sqrt{n}\}$ for which*

$$\sqrt{n}(\hat{\beta} - \beta_0) \longrightarrow_{\mathcal{L}} N\left(0, \frac{1}{4f^2(0)}\Sigma^{-1}\Omega\Sigma^{-1}\right),$$

where ξ is a normal random vector with mean 0 and covariance matrix $\frac{1}{4f^2(0)}\Sigma^{-1}\Omega\Sigma^{-1}$.

Notice that the lack of convexity for the objective function $W_n(\beta)$ complicates the search for its minimizer. As in Davis and Dunsmuir (1997), $W_n(\beta)$ may be linearized in a neighborhood of a good initial estimate $\hat{\beta}^0$ as follows

$$\tilde{W}_n(\beta) = \sum_{t=u+1}^n \tilde{w}_t |\varepsilon_t(\hat{\beta}^0) - A'_t(\hat{\beta}^0)(\beta - \hat{\beta}^0)|.$$

The resulting estimator $\tilde{\beta} = \arg \min_{\beta} \tilde{W}_n(\beta)$ shares the same asymptotic property as the local WLAD. See the theorem below.

Theorem 2. *Let conditions A1 – A3 hold. Then*

$$\sqrt{n}(\tilde{\beta} - \beta_0) \longrightarrow_{\mathcal{L}} N\left(0, \frac{1}{4f^2(0)}\Sigma^{-1}\Omega\Sigma^{-1}\right),$$

provided that $\hat{\beta}^0 = \beta_0 + O_p(n^{-1/2})$.

Remark 3. Although we only deal with the weight function defined in (2.4) explicitly, the theorem above holds for general weight function $g_t \equiv g(y_{t-1}, y_{t-2}, \dots)$ provided

$$E\{(g_t + g_t^2)(\xi^2 + \xi^3)\} < \infty, \quad E\{(g_t + g_t^2)(\xi_t^2 + \xi_t^3 + \xi_t^4)\} < \infty$$

where $\xi = \sum_{i=0}^{\infty} r^i |y_{-i}|$, $\xi_t = C_0 \sum_{i=t}^{\infty} r^i |y_{t-i}|$, $0 < r < 1$ and $C_0 > 0$ are constants. For example, we may use the weights of more general form

$$g_t = \left(1 + \sum_{k=1}^{t-1} k^{-\alpha} (\log k)^d |y_{t-k}|\right)^{-\gamma}, \quad \alpha > 2, \quad \gamma \geq 2, \quad d \geq 0. \quad (2.8)$$

The numerical examples in section 3 indicates that the accuracy of the WLAD is not sensitive with respect to the value of α . However the choice of $\gamma = 2$ typically leads a better estimator than those with $\gamma > 2$, at least for model (4.1). Furthermore, it seems that $d = 2, 3, 4$ behave almost equally well. However it remains an open question how to choose a weight function in general such that the resulting estimator is of certain optimality.

2.3 A Wald test for linear hypotheses

The asymptotic normality of the estimator $\hat{\beta}$ stated in Theorem 1 above facilitates the inference for model (1.1). For example, we may consider a general form of linear null hypothesis

$$H_0 : \Gamma\beta_0 = \kappa,$$

where Γ is a $s \times (p + q)$ constant matrix with rank s , and κ is $s \times 1$ constant vector. A Wald test statistic may be defined as

$$Z_n = (\Gamma\hat{\beta} - \kappa)' \left\{ \Gamma \frac{1}{4n\tilde{f}^2(0)} \hat{\Sigma}^{-1} \hat{\Omega} \hat{\Sigma}^{-1} \Gamma' \right\}^{-1} (\Gamma\hat{\beta} - \kappa)$$

and we reject H_0 for large values of Z_n . In the above expression,

$$\hat{\Sigma} = \frac{1}{n-u} \sum_{t=u+1}^n \tilde{w}_t \hat{Q}_t \hat{Q}_t', \quad \hat{\Omega} = \frac{1}{n-u} \sum_{t=u+1}^n (\tilde{w}_t^2 \hat{Q}_t \hat{Q}_t'), \quad (2.9)$$

\hat{Q}_t is defined in the same manner as Q_t but with β_0 replaced by $\hat{\beta}$, ε_t replaced by $\varepsilon_t(\hat{\beta})$ and $y_t = 0$ for all $t \leq 0$; see (2.6) and (2.7), and $\tilde{f}(0)$ is an estimate for $f(0)$ defined as

$$\tilde{f}(0) = \frac{1}{b_n} \sum_{t=u+1}^n \tilde{w}_t K\left(\frac{\varepsilon_t(\hat{\beta})}{b_n}\right) / \sum_{t=u+1}^n \tilde{w}_t, \quad (2.10)$$

where $K(\cdot)$ is a kernel function on R , $b_n > 0$ is a bandwidth. The theorem below shows that the asymptotic null-distribution of Z_n is χ_s^2 . It in fact still holds if $\hat{\beta}$ in the definition of Z_n is replaced by $\tilde{\beta}$.

Theorem 3. *Suppose conditions A1 – A3 hold. Let kernel function K be bounded, Lipschitz continuous and of finite first moment. Let $b_n \rightarrow 0$ and $nb_n^4 \rightarrow \infty$ as $n \rightarrow \infty$. Then $Z_n \rightarrow_{\mathcal{L}} \chi_s^2$ under H_0 .*

3 PROOFS

We use the same notation as in section 2. For any fixed $v \in R^{p+q}$, put

$$\begin{aligned} S_n(v) &= \sum_{t=u+1}^n w_t (|\varepsilon_t(\beta_0 + n^{-1/2}v)| - |\varepsilon_t(\beta_0)|), \\ S_n^+(v) &= \sum_{t=u+1}^n w_t (|\varepsilon_t - n^{-1/2}Q_t'v| - |\varepsilon_t|), \\ S_n^*(v) &= \sum_{t=u+1}^n w_t (|\varepsilon_t(\beta_0) - n^{-1/2}A_t'(\beta_0)v| - |\varepsilon_t(\beta_0)|). \end{aligned}$$

We linearize $\varepsilon_t(\beta)$ around β_0 , i.e. $\varepsilon_t(\beta)$ is approximated by

$$\varepsilon_t(\beta_0) - A'_t(\beta_0)(\beta - \beta_0),$$

where $A_t(\beta)$ is defined in (2.5).

We denote by “ $\rightarrow_{\mathcal{L}}$ ” the convergence in distribution, “ \xrightarrow{P} ” the convergence in probability. Let $C(R^s)$ be the space of the real-valued continuous functions on R^s (Rudin, 1991). For probability measures P_n and P on $C(R^s)$, we say that P_n converges weakly to P in $C(R^s)$ if $\int f dP_n \rightarrow \int f dP$ for any bounded and continuous function f defined on $C(R^s)$. For random functions S_n, S defined on $C(R^s)$, $S_n \rightarrow_{\mathcal{L}} S$ if the distribution of S_n converges weakly to that of S in $C(R^s)$ (Billingsley, 1999). C denotes a positive constant which may be different at different places.

3.1 Proof of Theorem 1

Before we prove Theorem 1, we first introduce a proposition which is of independent interest. Its proof is divided into several lemmas. We always assume that conditions A1 – A3 hold.

Proposition 2. *As $n \rightarrow \infty$, it holds that $T_n(v) \rightarrow_{\mathcal{L}} T(v)$ on $C(R^{p+q})$, where $T(v) = f(0)v'\Sigma v + v'N$, and N denotes a $N(0, \Omega)$ random vector.*

Lemma 1. *It holds that $|\varepsilon_t - \varepsilon_t(\beta_0)| \leq \xi_t$, and $|A_{t,i}(\beta_0) - Q_{t,i}| \leq \xi_t$ for $i = 1, \dots, p + q$, where $\xi_t = C_0 \sum_{j=t}^{\infty} r^j |y_{t-j}|$, $0 < r < 1$, C_0 is a positive constant, and $Q_{t,i}$ is the i -th component of Q_t .*

Proof. See page 265-268 in Brockwell and Davis (1991). □

Lemma 2. $S_n^+(v) \rightarrow_{\mathcal{L}} T(v)$ on $C(R^{p+q})$.

Proof. We first prove the convergence for any fixed v . Using the identity

$$|z - y| - |z| = -y \operatorname{sgn}(z) + 2(y - z)\{I(0 < z < y) - I(y < z < 0)\},$$

which holds for $z \neq 0$, we have

$$\begin{aligned} S_n^+(v) &= -n^{-1/2} \sum_{t=u+1}^n w_t Q'_t v \operatorname{sgn}(\varepsilon_t) \\ &\quad + 2 \sum_{t=u+1}^n w_t (n^{-1/2} Q'_t v - \varepsilon_t) [I(0 < \varepsilon_t < n^{-1/2} Q'_t v) - I(n^{-1/2} Q'_t v < \varepsilon_t < 0)] \\ &=: A_n + B_n. \end{aligned}$$

Notice that, by Lemma 1, we have $|Q_{t,i}| \leq C \sum_{j=1}^{\infty} r^j |y_{t-j}|$ and

$$|w_t^{1/2} Q_{t,i}| \leq \frac{C \sum_{j=1}^{\infty} r^j |y_{t-j}|}{1 + \sum_{k=1}^{\infty} k^{-\alpha} |y_{t-k}|} \leq C \sum_{k=1}^{\infty} k^{\alpha} r^k. \quad (3.1)$$

Then, $Ew_t^2(Q'_t v)^2 < +\infty$. But, from conditions A1 and A2, $\{w_t Q'_t v \text{sgn}(\varepsilon_t)\}$ is a stationary martingale difference sequence. Therefore, applying a martingale central limit theorem (Hall and Heyde (1980)), we obtain $A_n \rightarrow_{\mathcal{L}} v'N$.

For B_n , let

$$W_{nt} = w_t(n^{-1/2}Q'_t v - \varepsilon_t)I(0 < \varepsilon_t < n^{-1/2}Q'_t v)$$

and $\mathcal{F}_{t-1} = \sigma(\varepsilon_j, j \leq t-1)$. Then

$$\begin{aligned} nEW_{nt}^2 &= nE(E(W_{nt}^2|\mathcal{F}_{t-1})) \\ &= nE(w_t^2[\int_0^{n^{-1/2}Q'_t v} (n^{-1/2}Q'_t v - z)^2(f(z) - f(0))dz \\ &\quad + \int_0^{n^{-1/2}Q'_t v} (n^{-1/2}Q'_t v - z)^2 f(0)dz]) \\ &\leq nE(w_t^2 B_2 n^{-2}(Q'_t v)^4 + w_t^2 B_1 n^{-3/2}(Q'_t v)^3). \end{aligned}$$

Similarly to (3.1), we can obtain

$$Ew_t^2(Q'_t v)^4 < +\infty, \quad Ew_t^2(Q'_t v)^3 < +\infty.$$

Therefore, we have proved that

$$\limsup_{n \rightarrow \infty} nEW_{nt}^2 = 0. \quad (3.2)$$

On the other hand, on the set $\{Q'_t v > 0\}$, we may show that

$$\sum_{t=u+1}^n E(W_{nt}|\mathcal{F}_{t-1}) \rightarrow \frac{f(0)}{2} E[w_t(Q'_t v)^2 I(Q'_t v > 0)]$$

and

$$\text{Var}\left(\sum_{t=u+1}^n (W_{nt} - E(W_{nt}|\mathcal{F}_{t-1}))\right) \rightarrow 0,$$

see Davis and Dunsmuir (1997). Therefore,

$$\sum_{t=u+1}^n W_{nt} \rightarrow \frac{f(0)}{2} E[w_t(Q'_t v)^2 I(Q'_t v > 0)]. \quad (3.3)$$

Using the same argument for the second indicator in the summands of B_n , we obtain that

$$B_n \xrightarrow{P} f(0)v'\Sigma v. \quad (3.4)$$

So that the finite dimensional distributions of S_n^+ converge to those of T . But since S_n^+ has convex sample paths, this implies that the convergence is in fact on $C(R^{p+q})$ (see the proof of Proposition 1 in Davis and Dunsmuir (1997)). \square

Lemma 3. $S_n^*(v) - S_n^+(v) \xrightarrow{P} 0$ uniformly on compact sets.

Proof. Notice that

$$\begin{aligned} & S_n^*(v) - S_n^+(v) \\ = & \sum_{t=u+1}^n w_t [(|\varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v| - |\varepsilon_t(\beta_0)|) - (|\varepsilon_t - n^{-1/2}Q'_t v| - |\varepsilon_t|)] \\ = & -n^{-1/2} \sum_{t=u+1}^n w_t A'_t(\beta_0)v \operatorname{sgn}(\varepsilon_t(\beta_0)) + n^{-1/2} \sum_{t=u+1}^n w_t Q'_t v \operatorname{sgn}(\varepsilon_t) \\ & + 2 \sum_{t=u+1}^n w_t (n^{-1/2}A'_t(\beta_0)v - \varepsilon_t(\beta_0)) [I(0 < \varepsilon_t(\beta_0) < n^{-1/2}A'_t(\beta_0)v) \\ & - I(n^{-1/2}A'_t(\beta_0)v < \varepsilon_t(\beta_0) < 0)] - 2 \sum_{t=u+1}^n w_t (n^{-1/2}Q'_t v - \varepsilon_t) \\ & [I(0 < \varepsilon_t < n^{-1/2}Q'_t v) - I(n^{-1/2}Q'_t v < \varepsilon_t < 0)]. \end{aligned}$$

First, we consider

$$\begin{aligned} \Lambda_1 & = n^{-1/2} \sum_{t=u+1}^n w_t [Q'_t v \operatorname{sgn}(\varepsilon_t) - A'_t(\beta_0)v \operatorname{sgn}(\varepsilon_t(\beta_0))] \\ & = n^{-1/2} \sum_{t=u+1}^n w_t Q'_t v (\operatorname{sgn}(\varepsilon_t) - \operatorname{sgn}(\varepsilon_t(\beta_0))) \\ & \quad + n^{-1/2} \sum_{t=u+1}^n w_t (Q'_t v - A'_t(\beta_0)v) \operatorname{sgn}(\varepsilon_t(\beta_0)) \\ & =: K_1 + K_2. \end{aligned}$$

By Lemma 1 and the proof of (3.1), we know

$$|w_t(Q'_t v - A'_t(\beta_0)v)| \leq C\|v\|\xi_t.$$

Then

$$K_2 \leq n^{-1/2} C \|v\| \sum_{t=u+1}^n \sum_{j=t}^{\infty} r^j |y_{t-j}| = n^{-1/2} C \|v\| \sum_{t=u+1}^n r^t \sum_{h=0}^{\infty} r^h |y_{-h}| \xrightarrow{P} 0.$$

uniformly on compact sets. For K_1 , we have

$$\begin{aligned} E|K_1| &\leq n^{-1/2} \sum_{t=u+1}^n E(E(w_t | Q'_t v) | \text{sgn}(\varepsilon_t) - \text{sgn}(\varepsilon_t(\beta_0)) | \mathcal{F}_{t-1}) \\ &= 2n^{-1/2} \sum_{t=u+1}^n E(w_t | Q'_t v) P(\{\varepsilon_t > 0, \varepsilon_t(\beta_0) < 0\} \cup \{\varepsilon_t < 0, \varepsilon_t(\beta_0) > 0\} | \mathcal{F}_{t-1}) \\ &\leq 2n^{-1/2} \sum_{t=u+1}^n E(w_t | Q'_t v) P(\{0 < \varepsilon_t < \xi_t\} \cup \{-\xi_t < \varepsilon_t < 0\} | \mathcal{F}_{t-1}) \\ &= 2n^{-1/2} \sum_{t=u+1}^n [E(w_t | Q'_t v) \int_{-\xi_t}^{\xi_t} f(x) dx] \leq 4B_1 n^{-1/2} \sum_{t=u+1}^n E(w_t^{1/2} | Q'_t v) \cdot w_t^{1/2} \xi_t \\ &\leq 4CB_1 \|v\| n^{-1/2} \sum_{t=u+1}^n \sum_{h=0}^{\infty} (t+h)^\alpha r^{t+h} \\ &\leq 4CB_1 \|v\| n^{-1/2} \left[\sum_{t=u+1}^n t^\alpha r^t \sum_{h=0}^{\infty} r^h + \sum_{t=u+1}^n r^t \sum_{h=0}^{\infty} h^\alpha r^h \right] \rightarrow 0 \end{aligned}$$

uniformly on compact sets, because we have the facts

$$w_t^{1/2} |Q'_t v| \leq C \|v\|, \quad w_t^{1/2} \xi_t \leq \sum_{j=t}^{\infty} r^j j^\alpha \quad \text{and} \quad (t+h)^\alpha \leq 2^\alpha (t^\alpha + h^\alpha).$$

Therefore, $\Lambda_1 \xrightarrow{P} 0$ uniformly on compact sets as $n \rightarrow \infty$.

Now we consider

$$\begin{aligned} \Lambda_2 &= \sum_{t=u+1}^n w_t (n^{-1/2} A'_t(\beta_0) v - \varepsilon_t(\beta_0)) I(0 < \varepsilon_t(\beta_0) < n^{-1/2} A'_t(\beta_0) v) \\ &\quad - \sum_{t=u+1}^n w_t (n^{-1/2} Q'_t v - \varepsilon_t) I(0 < \varepsilon_t < n^{-1/2} Q'_t v) \\ &= \sum_{t=u+1}^n w_t (n^{-1/2} A'_t(\beta_0) v - \varepsilon_t(\beta_0) - n^{-1/2} Q'_t v + \varepsilon_t) I(0 < \varepsilon_t(\beta_0) < n^{-1/2} A'_t(\beta_0) v) \\ &\quad + \sum_{t=u+1}^n w_t (n^{-1/2} Q'_t v - \varepsilon_t) [I(0 < \varepsilon_t(\beta_0) < n^{-1/2} Q'_t v) - I(0 < \varepsilon_t < n^{-1/2} Q'_t v)] \\ &\quad + \sum_{t=u+1}^n w_t (n^{-1/2} Q'_t v - \varepsilon_t) [I(0 < \varepsilon_t(\beta_0) < n^{-1/2} A'_t(\beta_0) v) - I(0 < \varepsilon_t(\beta_0) < n^{-1/2} Q'_t v)] \\ &= D_1 + D_2 + D_3 \end{aligned}$$

By the similar way to the proof for K_2 , we can obtain that $D_1 \xrightarrow{P} 0$ uniformly on compact sets.

On the other hand, denote $\eta_t = \varepsilon_t(\beta_0) - \varepsilon_t$. It can be verified that

$$\begin{aligned} & |I(0 < \varepsilon_t(\beta_0) < n^{-1/2}Q'_t v) - I(0 < \varepsilon_t < n^{-1/2}Q'_t v)| \\ & \leq (I(-\eta_t < \varepsilon_t < 0)) + I(n^{-1/2}Q'_t v - \eta_t < \varepsilon_t < n^{-1/2}Q'_t v)I(\eta_t > 0) \\ & \quad + (I(0 < \varepsilon_t < -\eta_t)) + I(n^{-1/2}Q'_t v < \varepsilon_t < n^{-1/2}Q'_t v - \eta_t)I(\eta_t < 0). \end{aligned}$$

Then from condition A3 and by the similar way to the proof of Lemma 2, we have

$$\begin{aligned} E|D_2| & \leq \sum_{t=u+1}^n E\{w_t I(Q'_t v > 0) [I(\eta_t > 0) \left(\int_{-\eta_t}^0 + \int_{n^{-1/2}Q'_t v - \eta_t}^{n^{-1/2}Q'_t v} \right) |n^{-1/2}Q'_t v - x| f(x) dx \\ & \quad + I(\eta_t < 0) \left(\int_0^{-\eta_t} + \int_{n^{-1/2}Q'_t v}^{n^{-1/2}Q'_t v - \eta_t} \right) |n^{-1/2}Q'_t v - x| f(x) dx]\} \\ & \leq C \sum_{t=u+1}^n E\{w_t (|n^{-1/2}Q'_t v| \xi_t + \xi_t^2)\} \\ & = C n^{-1/2} \sum_{t=u+1}^n E\{w_t^{1/2} |Q'_t v| \cdot w_t^{1/2} \xi_t\} + C \sum_{t=u+1}^n E\{w_t^{1/2} \xi_t\}^2 \\ & = D_{21} + D_{22} \end{aligned}$$

By the same method as for K_1 , we can obtain that $D_{21} \rightarrow 0$. On the other hand,

$$\begin{aligned} D_{22} & \leq C \sum_{t=u+1}^n \left(\sum_{h=0}^{\infty} r^{h+t} (h+t)^\alpha \right)^2 \leq C \sum_{t=u+1}^n r^{2t} \left(\sum_{h=0}^{\infty} r^h h^\alpha + \sum_{h=0}^{\infty} r^h t^\alpha \right)^2 \\ & \leq C \sum_{t=u+1}^n r^{2t} \left(\sum_{h=0}^{\infty} r^h h^\alpha \right)^2 + \sum_{t=u+1}^n r^{2t} \left(\sum_{h=0}^{\infty} r^h t^\alpha \right)^2 \rightarrow 0. \end{aligned}$$

Therefore, we have

$$E|D_2| \longrightarrow 0. \tag{3.5}$$

And we can obtain $E|D_3| \longrightarrow 0$ similarly to (3.5). Hence $\Lambda_2 \xrightarrow{P} 0$ uniformly on compact sets.

By the same way for Λ_2 , we have

$$\begin{aligned} \Lambda_3 & = \sum_{t=u+1}^n w_t (n^{-1/2}A'_t(\beta_0)v - \varepsilon_t(\beta_0)) I(n^{-1/2}A'_t(\beta_0)v < \varepsilon_t(\beta_0) < 0) \\ & \quad - \sum_{t=u+1}^n w_t (n^{-1/2}Q'_t v - \varepsilon_t) I(n^{-1/2}Q'_t v < \varepsilon_t < 0) \xrightarrow{P} 0 \end{aligned}$$

uniformly on compact sets. This completes the proof of this lemma. \square

Lemma 4. $S_n(v) - T_n(v) \xrightarrow{P} 0$ uniformly on compact sets.

Proof. Notice that, we have the fact that under conditions A1 and A2, for any given positive number M , there exist positive constants $C > 0$ and $0 < r < 1$ such that for sufficiently large n

$$\sup_{\|v\| \leq M} |\varepsilon_t(\beta_0 + n^{-1/2}v)| \leq C \sum_{j=0}^{\infty} r^j |y_{t-j}|,$$

Now we have

$$\begin{aligned} \sup_{\|v\| \leq M} |S_n(v) - T_n(v)| &\leq \sup_{\|v\| \leq M} \sum_{t=u+1}^n |w_t - \tilde{w}_t| [|\varepsilon_t(\beta_0 + n^{-1/2}v)| + |\varepsilon_t(\beta_0)|] \\ &\leq C \sum_{t=u+1}^n \frac{\sum_{k=t}^{\infty} k^{-\alpha} |y_{t-k}| \sum_{j=0}^{\infty} r^j |y_{t-j}|}{1 + \sum_{k=1}^{\infty} k^{-\alpha} |y_{t-k}|} \leq C \sum_{t=u+1}^n \sum_{k=t}^{\infty} k^{-\alpha} |y_{t-k}| (\sum_{j=1}^{\infty} r^j j^\alpha + |y_t|) \\ &\leq C \sum_{t=u+1}^n |y_t| \sum_{h=0}^{\infty} (h+t)^{-\alpha} |y_{-h}| + C \sum_{t=u+1}^n \sum_{h=0}^{\infty} (h+t)^{-\alpha} |y_{-h}| \\ &\leq C \left(\sum_{t=u+1}^n t^{-\alpha/2} (|y_t| + 1) \right) \sum_{h=0}^{\infty} h^{-\alpha/2} |y_{-h}| \xrightarrow{P} 0 \end{aligned}$$

as $n \rightarrow \infty$, by conditions A2 and A3 and the fact $(h+t)^{-\alpha} \leq 2^{-\alpha} (ht)^{-\alpha/2}$. This completes the proof of Lemma 4. \square

Proof of Proposition 2. For $v \in R^{p+q}$, define

$$S_n^{**}(v) = \sum_{t=u+1}^n w_t (|\varepsilon_t(\beta_0) - n^{-1/2} A'_t(\beta_0)v - n^{-1} v' H_t(\beta_0)v| - |\varepsilon_t(\beta_0)|)$$

where

$$H_t(\beta) = -\frac{1}{2} \frac{\partial^2 \varepsilon_t(\beta)}{\partial \beta \partial \beta'} = (h_{t(i,j)})_{(p+q) \times (p+q)}.$$

From the definition of $\varepsilon_t(\beta)$, we have

$$\theta(B) \frac{\partial^2 \varepsilon_t(\beta)}{\partial \phi_i \partial \phi_j} = 0, \quad i, j = 1, \dots, p,$$

$$\theta(B) \frac{\partial^2 \varepsilon_t(\beta)}{\partial \phi_i \partial \theta_j} = A_{t-j,i}(\beta), \quad i = 1, \dots, p, j = 1, \dots, q,$$

$$\theta(B) \frac{\partial^2 \varepsilon_t(\beta)}{\partial \theta_i \partial \theta_j} = A_{t-j,i}(\beta) + A_{t-i,j}(\beta), \quad i, j = 1, \dots, q.$$

Similarly, replacing $A_{t-i,j}$ (or $A_{t-j,i}$) by $Q_{t-i,j}$ (or $Q_{t-j,i}$) in the above three equalities, we can define $X_t = (X_{t(i,j)})$. Then, $(h_{t(i,j)}(\beta_0))$ can be well approximated by $(X_{t(i,j)})$ such that

$$|X_{t(i,j)} - h_{t(i,j)}(\beta_0)| \leq \xi_t, \quad i, j = 1, \dots, p+q, \quad (3.6)$$

which can be proved by the same way as that for Lemma 1.

Notice that

$$\begin{aligned} & S_n^{**}(v) - S_n^*(v) \\ &= \sum_{t=u+1}^n w_t (|\varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v - n^{-1}v'H_t(\beta_0)v| - |\varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v|) \\ &= -\frac{1}{n} \sum_{t=u+1}^n w_t v' H_t(\beta_0) v \operatorname{sgn}(\varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v) \\ &\quad + 2 \sum_{t=u+1}^n w_t (n^{-1}v'H_t(\beta_0)v - \varepsilon_t(\beta_0) + n^{-1/2}A'_t(\beta_0)v) \\ &\quad [I(0 < \varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v < n^{-1}v'H_t(\beta_0)v) \\ &\quad - I(n^{-1}v'H_t(\beta_0)v < \varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v < 0)] \\ &= T_1 + T_2. \end{aligned}$$

Notice that $-\frac{1}{n} \sum_{t=u+1}^n w_t v' X_t v \operatorname{sgn}(\varepsilon_t) \rightarrow 0$, we obtain

$$T_1 = -\frac{1}{n} \sum_{t=u+1}^n w_t v' X_t v [\operatorname{sgn}(\varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v) - \operatorname{sgn}(\varepsilon_t)] + o_p(1)$$

by (3.6). Using the same argument of $E|K_1| \rightarrow 0$ in Lemma 3, we have $T_1 \xrightarrow{P} 0$. Similarly, we have

$$\begin{aligned} T_{21} &= \sum_{t=u+1}^n w_t (n^{-1}v'H_t(\beta_0)v - \varepsilon_t(\beta_0) + n^{-1/2}A'_t(\beta_0)v) \\ &\quad I(0 < \varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v < n^{-1}v'H_t(\beta_0)v) \\ &= \sum_{t=u+1}^n w_t (n^{-1}v'X_t v - \varepsilon_t + n^{-1/2}Q'_t v) I(0 < \varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v < n^{-1}v'H_t(\beta_0)v) \\ &\quad + o_p(1). \end{aligned}$$

Let

$$T_{21}^{(1)} = \sum_{t=u+1}^n w_t (n^{-1}v'X_t v - \varepsilon_t + n^{-1/2}Q'_t v) I(0 < \varepsilon_t - n^{-1/2}Q'_t v < n^{-1}v'X_t v)$$

and

$$T_{21}^{(2)} = \sum_{t=u+1}^n w_t (n^{-1}v'X_tv - \varepsilon_t + n^{-1/2}Q'_tv) [I(0 < \varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v < n^{-1}v'H_t(\beta_0)v) - I(0 < \varepsilon_t - n^{-1/2}Q'_tv < n^{-1}v'X_tv)].$$

Using the same way as the proof of (3.3) and (3.5) for $T_{21}^{(1)}$ and $T_{21}^{(2)}$ respectively, we can obtain that $T_{21} \xrightarrow{P} 0$. The same result holds for the rest term of T_2 . Thus, it follows that $S_n^{**}(v) - S_n^*(v) \xrightarrow{P} 0$ uniformly on compact sets.

But,

$$\begin{aligned} & |S_n^{**}(v) - S_n(v)| \\ &= \left| \sum_{t=u+1}^n w_t (|\varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v - n^{-1}v'H_t(\beta_0)v| - |\varepsilon_t(\beta_0 + n^{-1/2}v)|) \right| \\ &\leq \frac{1}{n} \sum_{t=1}^n w_t |v'(H_t(\beta_n^*) - H_t(\beta_0))v| \end{aligned}$$

where β_n^* is between β_0 and $\beta_0 + n^{-1/2}v$. Then, $S_n^{**}(v) - S_n(v) \xrightarrow{P} 0$ uniformly on compact sets. Therefore, combining Lemma 3 and Lemma 4, we have $T_n(v) - S_n^+(v) \xrightarrow{P} 0$ uniformly on compact sets.

By Lemma 2, we obtain that

$$T_n(v) \rightarrow_{\mathcal{L}} T(v) \quad \text{on } C(R^{p+q})$$

as $n \rightarrow \infty$. This completes the proof.

Proof of Theorem 1. Note that the limit process in Proposition 2 has convex sample paths and a unique minimizer

$$\xi = -(1/[2f(0)])\Sigma^{-1}N.$$

Denote P_n and P be the probability measures on $C(R^{p+q})$ induced by T_n and T respectively. By Skorokhod's representation theorem (cf. Pollard (1984), page 71-73), there exists a probability space $(\Omega^*, \mathcal{F}^*, P^*)$ with processes T^* and T_n^* having the distributions P and P_n respectively such that $T_n^* \xrightarrow{a.s.} T^*$. Hence, there exists a subset Ω' of Ω^* with $P^*(\Omega') = 1$ such that for any

$\omega \in \Omega'$,

$$\sup_{v \in K} |T_n^*(v, \omega) - T^*(v, \omega)| \rightarrow 0. \quad (3.7)$$

holds for any compact set K . Denote the minimizer of T^* by ξ^* . Then ξ^* has the same distribution as ξ . For any given positive random variable M , let

$$\xi_n^*(\omega) = \arg \min_{\|v - \xi^*(\omega)\| \leq M(\omega)} T_n^*(v, \omega).$$

Now we show that $\xi_n^*(\omega) \rightarrow \xi^*(\omega)$. Suppose $\xi_n^*(\omega) \rightarrow \xi^*(\omega)$ doesn't hold, then there is a subsequence n' such that $\xi_{n'}^*(\omega) \rightarrow \xi'(\omega) \neq \xi^*(\omega)$ and we know $\|\xi'(\omega) - \xi^*(\omega)\| \leq M(\omega)$. From the definition of $\xi_n^*(\omega)$, we have

$$T_n^*(\xi^*(\omega)) - T_n^*(\xi_{n'}^*(\omega)) \geq 0.$$

But, on the other hand,

$$\begin{aligned} & T_{n'}^*(\xi^*(\omega)) - T_{n'}^*(\xi_{n'}^*(\omega)) \\ &= T_{n'}^*(\xi^*(\omega)) - T^*(\xi^*(\omega)) + T^*(\xi^*(\omega)) - T^*(\xi_{n'}^*(\omega)) + T^*(\xi_{n'}^*(\omega)) - T_{n'}^*(\xi_{n'}^*(\omega)) \\ &= G_1 + G_2 + G_3. \end{aligned}$$

From (3.7), we can obtain that $G_1 \rightarrow 0$ and $G_3 \rightarrow 0$. Noticing that $T^*(v, \omega)$ is continuous, we have

$$G_2 \rightarrow T^*(\xi^*(\omega)) - T^*(\xi'(\omega)) < 0.$$

This is a contradiction. Therefore, $\xi_n^* \xrightarrow{a.s.} \xi^*$.

Define a sequence of local minimizers $\{\xi_n\}$ of T_n by

$$\xi_n(\omega) = \arg \min_{\|v - \xi(\omega)\| \leq M(\omega)} T_n(v, \omega).$$

Then, ξ_n converges in distribution to ξ . The proof is completed.

3.2 Proof of Theorem 2

We also use the substitution $v = \sqrt{n}(\beta - \beta_0)$. Then, minimizing $\tilde{W}_n(\beta)$ is equivalent to minimizing

$$\tilde{T}_n(v) = \sum_{t=u+1}^n \tilde{w}_t(|\varepsilon_t(\hat{\beta}^0) - A'_t(\hat{\beta}^0)(\beta_0 - \hat{\beta}^0 + n^{-1/2}v)| - |\varepsilon_t|).$$

Since $\tilde{T}_n(v)$ has convex sample paths, we only need to prove that $\tilde{T}_n(v) \rightarrow_{\mathcal{L}} T(v)$ on $C(R^{p+q})$ by Lemma 2.2 in Davis et. al. (1992).

Denote

$$\tilde{H}_t(\beta) = \frac{1}{2} \sum_{i,j=1}^{p+q} \frac{\partial^2 \varepsilon_t(\beta)}{\partial \beta_i \partial \beta_j}.$$

By Taylor expansion near $\hat{\beta}^0$, for any fixed $v \in R^{p+q}$, we have

$$\varepsilon_t(\beta_0 + n^{-1/2}v) = \varepsilon_t(\hat{\beta}^0) - A'_t(\hat{\beta}^0)(\beta_0 - \hat{\beta}^0 + n^{-1/2}v) + \tilde{H}_t(\beta_{1t})O_p(n^{-1}),$$

where β_{1t} lies between $\hat{\beta}^0$ and $\beta_0 + n^{-1/2}v$. We now have

$$\begin{aligned} & \varepsilon_t(\hat{\beta}^0) - A'_t(\hat{\beta}^0)(\beta_0 - \hat{\beta}^0 + n^{-1/2}v) \\ &= \varepsilon_t(\beta_0 + n^{-1/2}v) - \tilde{H}_t(\beta_{1t})O_p(n^{-1}) \\ &= \varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v + \tilde{H}_t(\beta_{2t})O_p(n^{-1}) - \tilde{H}_t(\beta_{1t})O_p(n^{-1}), \end{aligned}$$

where β_{2t} lies between β_0 and $\beta_0 + n^{-1/2}v$. Hence, by a similar way to the proof of Lemma 2 but replacing $n^{-1/2}Q'_t v$ by $n^{-1/2}Q'_t v - \tilde{H}_t(\beta_0)O_p(n^{-1})$, we have

$$\begin{aligned} \tilde{T}_n(v) &= \sum_{t=u+1}^n \tilde{w}_t [|\varepsilon_t(\beta_0) - n^{-1/2}A'_t(\beta_0)v + \tilde{H}_t(\beta_{2t})O_p(n^{-1}) - \tilde{H}_t(\beta_{1t})O_p(n^{-1})| - |\varepsilon_t|] \\ &= \sum_{t=u+1}^n \tilde{w}_t (|\varepsilon_t - n^{-1/2}Q'_t v + \tilde{H}_t(\beta_0)O_p(n^{-1})| - |\varepsilon_t|) + o_p(1) \rightarrow_{\mathcal{L}} T(v). \end{aligned}$$

3.3 Proof of Theorem 3

Based on Theorem 1, Theorem 3 follows immediately from Lemmas 5 & 6 below. We assume that the conditions of Theorem 3 holds.

Lemma 5. $\hat{\Sigma} \xrightarrow{P} \Sigma$ and $\hat{\Omega} \xrightarrow{P} \Omega$.

Proof. Denote

$$\hat{\theta}(z) = 1 + \hat{\theta}_1 z + \cdots + \hat{\theta}_q z^q, \quad \hat{\phi}(z) = 1 - \hat{\phi}_1 z - \cdots - \hat{\phi}_p z^p.$$

Then

$$\hat{Q}_t = (\hat{U}_{t-1}, \cdots, \hat{U}_{t-p}, \hat{V}_{t-1}, \cdots, \hat{V}_{t-q})'$$

where \hat{U}_t and \hat{V}_t are determined by

$$\hat{\phi}(B)\hat{U}_t = \varepsilon_t(\hat{\beta}), \quad \hat{\theta}(B)\hat{V}_t = \varepsilon_t(\hat{\beta}).$$

Note that, by the definition of $\varepsilon_t(\beta)$, we have $\hat{\phi}(B)y_t = \hat{\theta}(B)\varepsilon_t(\hat{\beta})$ where $y_t = 0$ for $t \leq 0$. Hence,

$$\hat{U}_t = \hat{\theta}^{-1}(B)y_t, \quad \hat{V}_t = \hat{\theta}^{-2}(B)\hat{\phi}(B)y_t, \quad (y_t = 0 \text{ for } t \leq 0).$$

Let

$$II_1 = \frac{1}{n-u} \sum_{t=u+1}^n \tilde{w}_t \hat{Q}_t \hat{Q}'_t - \frac{1}{n-u} \sum_{t=u+1}^n w_t Q_t Q'_t, \quad II_2 = \frac{1}{n-u} \sum_{t=u+1}^n w_t Q_t Q'_t.$$

Then, $\hat{\Sigma} = II_1 + II_2$. Obviously, $II_2 \rightarrow \Sigma$ by the ergodic theorem. For II_1 , we first define a vector-valued function $Q_t(\beta) = (U_{t-1}(\beta), \dots, U_{t-p}(\beta), V_{t-1}(\beta), \dots, V_{t-q}(\beta))'$ and its components are determined as follows

$$\phi(B)U_t(\beta) = \varepsilon_t, \quad \theta(B)V_t(\beta) = \varepsilon_t.$$

We denote $\tilde{Q}_t = Q_t(\hat{\beta})$ and divide II_1 into three terms as follows

$$\begin{aligned} II_1 &= \frac{1}{n-u} \sum_{t=u+1}^n (\tilde{w}_t - w_t) \hat{Q}_t \hat{Q}'_t + \frac{1}{n-u} \sum_{t=u+1}^n w_t (\hat{Q}_t \hat{Q}'_t - \tilde{Q}_t \tilde{Q}'_t) \\ &\quad + \frac{1}{n-u} \sum_{t=u+1}^n w_t (\tilde{Q}_t \tilde{Q}'_t - Q_t Q'_t) \\ &= J_1 + J_2 + J_3 \end{aligned}$$

It is easy to obtain that $J_1 \xrightarrow{P} 0$ by Lemma 1 and the definition of \hat{Q}_t . Notice that the $(1,1)th$ element of J_2

$$J_2^{(1,1)} = \frac{1}{n-u} \sum_{t=u+1}^n w_t (\hat{U}_{t-1} - U_{t-1}(\hat{\beta})) (\hat{U}_{t-1} + U_{t-1}(\hat{\beta}))$$

and

$$w_t^{1/2} |\hat{U}_{t-1} - U_{t-1}(\hat{\beta})| \leq w_t^{1/2} \xi_t \leq C, \quad w_t^{1/2} |\hat{U}_{t-1} + U_{t-1}(\hat{\beta})| \leq C$$

for some positive constant C and ξ_t defined in Lemma 1. It follows that $J_2^{(1,1)} \xrightarrow{P} 0$, and similarly we can prove that the rest elements of J_2 converges to zero in probability. Hence $J_2 \xrightarrow{P} 0$. For J_3 , noticing that $Q_t = Q_t(\beta_0)$, we have

$$\tilde{Q}_t - Q_t = Q_t(\hat{\beta}) - Q_t(\beta_0) = \frac{\partial Q_t(\beta^*)}{\partial \beta'} (\hat{\beta} - \beta_0).$$

where β^* lies between $\hat{\beta}$ and β_0 . Noticing that $w_t^{1/2} \|\frac{\partial Q_t(\beta^*)}{\partial \beta'}\| \leq C$, $w_t^{1/2} \|Q_t\| \leq C$ and $w_t^{1/2} \|\tilde{Q}_t\| \leq C$, we obtain that $J_3 \xrightarrow{P} 0$. Now it has been proved that $\hat{\Sigma} \xrightarrow{P} \Sigma$.

By the same way we can prove $\hat{\Omega} \xrightarrow{P} \Omega$. □

Lemma 6. *It holds that $\hat{f}(0) \xrightarrow{P} f(0)$.*

Proof. Define

$$\hat{f}(0) = \frac{1}{\hat{\sigma}_w b_n (n-u)} \sum_{t=u+1}^n w_t K\left(\frac{\varepsilon_t(\hat{\beta})}{b_n}\right)$$

where $\hat{\sigma}_w = (n-u)^{-1} \sum_{t=u+1}^n w_t$, then

$$\begin{aligned} |\hat{f}(0) - \tilde{f}(0)| &\leq \left| \frac{1}{\hat{\sigma}_{\tilde{w}} b_n (n-u)} \sum_{t=u+1}^n (\tilde{w}_t - w_t) K\left(\frac{\varepsilon_t(\hat{\beta})}{b_n}\right) \right| \\ &\quad + \left| \frac{\sigma_{\tilde{w}} - \hat{\sigma}_w}{\hat{\sigma}_{\tilde{w}} \hat{\sigma}_w b_n (n-u)} \right| \sum_{t=u+1}^n w_t K\left(\frac{\varepsilon_t(\hat{\beta})}{b_n}\right) \\ &\leq \frac{C}{\hat{\sigma}_{\tilde{w}} b_n (n-u)} \sum_{t=u+1}^n \sum_{k=t}^{\infty} k^{-\alpha} |y_{t-k}| + O_p(1) \left| \frac{\sigma_{\tilde{w}} - \hat{\sigma}_w}{\hat{\sigma}_{\tilde{w}} \hat{\sigma}_w b_n} \right| \\ &\leq o_p(1) + O_p(1) \frac{1}{(n-u) b_n} \sum_{t=u+1}^n (\tilde{w}_t - w_t) \xrightarrow{P} 0 \end{aligned}$$

where $\hat{\sigma}_{\tilde{w}} = (n-u)^{-1} \sum_{t=u+1}^n \tilde{w}_t$. So we need to prove that

$$\hat{f}(0) \xrightarrow{P} f(0).$$

Notice that

$$|\hat{f}(0) - f(0)| \leq P_1 + P_2$$

where

$$P_1 = \left| \frac{1}{\hat{\sigma}_w b_n (n-u)} \sum_{t=u+1}^n w_t \left(K\left(\frac{\varepsilon_t(\hat{\beta})}{b_n}\right) - K\left(\frac{\varepsilon_t}{b_n}\right) \right) \right|$$

$$P_2 = \left| \frac{1}{\hat{\sigma}_w b_n (n-u)} \sum_{t=u+1}^n w_t K\left(\frac{\varepsilon_t}{b_n}\right) - f(0) \right|.$$

But, Lipschitz continuity of $K(x)$ insures that there exists a positive number L such that $|K(x) - K(y)| \leq L|x - y|$ for any x, y , and then, from Theorem 1 and the assumptions on b_n , we have

$$\begin{aligned}
P_1 &\leq \frac{L}{\hat{\sigma}_w b_n^2(n-u)} \sum_{t=u+1}^n w_t |\varepsilon_t(\hat{\beta}) - \varepsilon_t| \\
&\leq \frac{L}{\hat{\sigma}_w b_n^2(n-u)} \sum_{t=u+1}^n w_t (|\varepsilon_t(\hat{\beta}) - \varepsilon_t(\beta_0)| + |\varepsilon_t(\beta_0) - \varepsilon_t|) \\
&\leq \frac{L}{\hat{\sigma}_w b_n^2(n-u)} \sum_{t=u+1}^n w_t \|\hat{\beta} - \beta_0\| \|A_t(\beta^*)\| + \frac{C}{\hat{\sigma}_w b_n^2(n-u)} \sum_{t=u+1}^n \sum_{k=t}^{\infty} r^k |y_{t-k}| \\
&\leq \frac{C}{\hat{\sigma}_w b_n^2 \sqrt{n}} \sqrt{n} \|\hat{\beta} - \beta_0\| + \frac{C}{\hat{\sigma}_w b_n^2(n-u)} \sum_{t=u+1}^n r^t \sum_{h=0}^{\infty} r^h |y_{t-h}| \xrightarrow{P} 0
\end{aligned}$$

where β^* lies between $\hat{\beta}$ and β_0 . For P_2 , equivalently we need to prove that

$$I = \frac{1}{\hat{\sigma}_w b_n(n-u)} \sum_{t=u+1}^n w_t K\left(\frac{\varepsilon_t}{b_n}\right) \xrightarrow{P} f(0). \quad (3.8)$$

In fact, $I = \frac{1}{\hat{\sigma}_w} I_1 + f(0)$ where

$$I_1 = \frac{1}{b_n(n-u)} \sum_{t=1}^n w_t K\left(\frac{\varepsilon_t}{b_n}\right) - \frac{1}{n-u} \sum_{t=u+1}^n w_t f(0).$$

Notice that

$$\begin{aligned}
EI_1^2 &= \frac{1}{(n-u)^2} \sum_{t=u+1}^n E\left[w_t \left(\frac{1}{b_n} K\left(\frac{\varepsilon_t}{b_n}\right) - f(0)\right)\right]^2 \\
&\quad + \frac{2}{(n-u)^2} \sum_{u+1 \leq i < j \leq n} E\left[w_i w_j \left(\frac{1}{b_n} K\left(\frac{\varepsilon_i}{b_n}\right) - f(0)\right) \left(\frac{1}{b_n} K\left(\frac{\varepsilon_j}{b_n}\right) - f(0)\right)\right] \\
&= I_{11} + I_{12}.
\end{aligned}$$

But

$$\begin{aligned}
I_{11} &\leq \frac{2}{n-u} E\left[w_t^2 \left(K^2\left(\frac{\varepsilon_t}{b_n}\right) \frac{1}{b_n^2} + f^2(0)\right)\right] \\
&= \frac{2}{b_n(n-u)} \int_{-\infty}^{\infty} K^2(x) f(b_n x) dx E(w_t^2) + \frac{2f^2(0)}{n-u} E w_t^2 \rightarrow 0
\end{aligned}$$

by the assumptions on $K(x)$, and

$$\begin{aligned}
|I_{12}| &= \frac{2}{(n-u)^2} \left| \sum_{u+1 \leq i < j \leq n} E\left[w_i w_j \left(K\left(\frac{\varepsilon_i}{b_n}\right) \frac{1}{b_n} - f(0)\right)\right] \int_{-\infty}^{\infty} K(x) (f(b_n x) - f(0)) dx \right| \\
&\leq \int_{-\infty}^{\infty} K(x) |f(b_n x) - f(0)| dx E\left[\left(K\left(\frac{\varepsilon_i}{b_n}\right) \frac{1}{b_n} + f(0)\right)\right] \leq C b_n \rightarrow 0
\end{aligned}$$

by A3 and the assumptions on $K(x)$. Noticing that $\hat{\sigma}_w \rightarrow Ew_t$, we have (3.8). The proof of Lemma 6 is completed. \square

4 NUMERICAL PROPERTIES

We conducted a simulation study to illustrate the finite sample properties of the proposed WLADE in the five aspects, namely, (i) its accuracy, (ii) its sampling distribution, (iii) comparison with the unweighted LADE, the Whittle estimator (Mikosch et al. 1995), QMLE, and the Gauss-Newton estimator (Davis 1996), (iv) the selection of α , γ , and d in (2.8), and (v) the performance of the Wald test statistic Z_n .

We generated data from a simple ARMA(1,1) model

$$y_t = \phi_1 y_{t-1} + \varepsilon_t + \theta_1 \varepsilon_{t-1}, \quad (4.1)$$

with t_2 , Cauchy or $N(0,1)$ innovation distribution. Unless specified otherwise, we always set $u = 20$, $\alpha = 3$ and $d = 0$.

Tables 1 and 2 list the means and the standard deviations (SD) of the WLADE for ϕ_1 and θ_1 from the 1000 samples from model (4.1) with sample size $n = 200$ or 400 , and the true value $(\phi_1, \theta_1) = (0.4, 0.7)$, $(0.3, 0.5)$ or $(-0.5, -0.5)$. The estimates are very accurate in general, and the accuracy increases when the sample size increases from 200 to 400. We also included in the tables for asymptotic standard deviations (AD) derived from Theorem 1 with (Σ, Ω) replaced by their estimators in (2.9), and $f(0)$ replaced by (2.10) with the kernel $K(x) = e^{-x}/(1 + e^{-x})^2$ and the rule-of-thumb bandwidth $b_n = 1.06 \times n^{-1/5}$. The values of SD and AD are pretty close with each other.

To investigate the sampling distributions of the WLADE, we drew 16000 samples of size $n = 400$ from (4.1) with $(\phi_1, \theta_1) = (0.3, 0.5)$ and t_2 or Cauchy innovations. For each sample, the WLADE for both ϕ_1 and θ_1 were obtained. We divided $[-6, 6]$ into small intervals with equal length 0.2. For each small interval, we computed the (normalized) relative frequency for the occurrence of the event that the normalized WLADE falls into the interval. Those relative frequencies are plotted against the center of intervals in Figures 1 and 2. We superimposed the $N(0,1)$ density function in the figures, which is the limiting density of the normalized WLADE.

Even with sample size $n = 400$, the estimated values of the density functions match their asymptotic limits very well.

Figure 3 presents the boxplots of the average absolute error (AAE) $(|\hat{\phi}_1 - 0.3| + |\hat{\theta}_1 - 0.5|)/2$ for the unweighted LADE, the WLADE, the Whittle estimator, the QMLE, and the Gauss-Newton estimator from 1000 samples with sample size $n = 400$ drawn from (4.1). Here, we set $d = 2$. For the samples with heavy-tailed innovations, i.e. t_2 and Cauchy, the WLADE performed better than the Whittle estimator, QMLE, and Gauss-Newton estimator. In fact, the improvement from using the WLADE over the above three estimators is more pronounced when the tails are heavier (i.e. with Cauchy distribution). The Gauss-Newton estimator performed the best with Gaussian innovations. However it is noticeable that there was an efficiency loss due to the introduction of weights in the estimation, although such a loss was not significant at least in the setting used in our simulation.

Since the convergence rate of the WLADE is slower than the Gauss-Newton estimator (Mikosch et. al 1995, and Davis 1996), we compared the two estimators with large sample size n between 2000 to 8000. For each setting, 1000 samples were drawn from model (4.1) with Cauchy innovations. The parameters (ϕ_1, θ_1) were set at $(-0.3, 0.2)$, $(0.3, -0.5)$, $(-0.3, 0.5)$, $(0.1, -0.6)$, $(0.3, 0.5)$, $(0.6, 0.7)$, $(0.4, 0.6)$, $(-0.4, -0.6)$, $(-0.3, -0.4)$, and $u = \sqrt{n}$. It turned out that when $\phi_1\theta_1 > 0$, the WLADE performed better than the Gauss-Newton estimator for all $n \leq 8000$. However, when $\phi_1\theta_1 < 0$, the WLADE was better than the Gauss-Newton estimator only for $n \leq 3000$. The boxplots of the AAE for parameters $(\phi_1, \theta_1) = (-0.3, 0.2), (-0.3, -0.4), (0.6, 0.7)$ are displayed in Figure 4.

We also compared the WLADE using a general form of weights (2.8) with different α , γ , and d . To this end, we draw 1000 samples from model (4.1) with $(\phi_1, \theta_1) = (0.3, 0.5)$ and $n = 400$. Figure 5 presents the boxplots of the AAEs with t_2 , Cauchy and normal innovations. They suggest that the WLADE is fairly robust with respect to the value of α . However, the WLADE is more sensitive to γ . It is evident that we should choose γ as smaller as possible, i.e. $\gamma = 2$, which corresponds to the default weight function used in this paper. On the other hand, the performances with $d = 1, 2$ and 3 do not differ significantly.

Finally, we approximated the P -value of the Wald test proposed in section 2.3 for testing

AR(1) against ARMA(1,1) models by the relative frequency \hat{P} for the occurrence of the event that the AR(1) null hypothesis was rejected in a simulation with 1000 replications. The data were generated from (4.1) with $(\phi_1, \theta_1) = (0.3, 0)$ or $(\phi_1, \theta_1) = (0.3, 0.5)$, and innovation to be $t(2)$, Cauchy or $N(0, 1)$. We repeated the experiment with sample size equal to, respectively, 200, 400 and 600 for the nominal significance level between 0 and 0.1. Figure 6 plots the difference between \hat{P} and the nominal significance level against the nominal level with data generated from (4.1). With $(\phi_1, \theta_1) = (0.3, 0)$, the three panels on the left in Figure 6 indicates that the χ^2 -asymptotic approximation for the significance level is accurate, especially for $n = 400$ and 600. With $(\phi_1, \theta_1) = (0.3, 0.5)$, the three panels on the right illustrate that the test is powerful in detecting the departure from AR(1) hypothesis.

REFERENCES

- Adler, R.J., R. Feldaman & M. Taqqu (1997) *A User's Guide to Heavy Tails: Statistical techniques for analyzing heavy tailed distribution and processes*. Birkhauser.
- An, H.Z. & Z.G. Chen (1982) On convergence of LAD estimates in autoregression with infinite variance. *J. Multivariate Analysis* 12, 335-345.
- Billingsley, P. (1999) *Convergence of Probability Measures*. Wiley.
- Brockwell, P.J. & R.A. Davis (1991) *Time Series: Theory and Methods*. Springer-Verlag.
- Calder. M & R.A. Davis (1998) Inference for linear processes with stable noise. In R.J. Adler, R. Feldaman & M. Taqqu (eds.), *A Practical Guide to Heavy Tails*. Boston: Birkhauser. pp. 159-176.
- Davis, R.A. (1996) Gauss-Newton and M-estimation for ARMA processes with infinite variance. *Stochastic Processes and Their Applications* 63, 75-95.
- Davis, R.A. & W.T.M. Dunsmuir (1997) Least absolute deviation estimation for regression with ARMA errors. *Journal Theoretical Probability* 10, 481-497.

- Davis, R.A., K. Knight & J. Liu (1992) M-estimation for autoregressions with infinite variance. *Stochastic Processes and Their Applications* 40, 145-180.
- Davis, R.A. & S.I. Resnick (1985) Limit theory for moving averagers of random variables with regularly varying tail probabilities. *Annals of Probability* 13, 179-195.
- Davis, R.A. & S.I. Resnick (1986) Limit theory for the sample covariance and correlation functions of moving averages. *Annals of Statistics* 14, 533-558.
- Dunsmuir, W.T.M. & N.M. Spencer (1991) Strong consistency and asymptotic normality of L_1 estimates of the autoregressive moving-average model. *Journal of Time Series Analysis* 12, 95-104.
- Gross, S. & W.L. Steiger (1979) Least absolute deviation estimates in autoregressions with infinite variance. *Journal of Applied Probability* 16, 104-116.
- Hall, P.C. & C.C. Heyde (1980) *Martingale Limit Theory and Its Applications*. Academic Press.
- Hjort, N.L. & D. Pollard (1993) Asymptotics for minimisers of convex processes. Available at <http://www.stat.yale.edu/~pollard/Papers/convex.pdf>.
- Horvath, L. & F. Liese (2004) L_p -estimators in ARCH models. *Journal of Statistical Planning and Inference* 119, 277-309.
- Jansen, D. & C. De Vries (1991) On the frequency of large stock returns: putting booms and busts into perspective. *Review of Economics and Statistics* 73, 18-28.
- Koedijk, K.G., M.M.A. Schafgans & C.G. De Vries (1990) The tail index of exchange rate returns, *Journal of International Economics* 29, 93-108.
- Kokoszka, P.S. & M.S. Taqqu (1996) Parameter estimation for infinite variance fractional ARIMA. *Annals of Statistics* 24, 1880-1913.
- Ling, S. (2005) Self-weighted LAD estimation for infinite variance autoregressive models. *Journal of the Royal Statistical Society B* 67, 381-393.

- Mikosch, T., T. Gadrich & R.J. Adler (1995) Parameter estimation for ARMA models with infinite variance innovations. *Annals of Statistics* 23, 305-326.
- Pollard, D. (1984) *Convergence of Stochastic Processes*. Springer-Verlag.
- Resnick, S.I. (1997) Heavy tail modeling and teletraffic data (with discussions). *Annals of Statistics* 25, 1805-1869.
- Rudin, W. (1991) *Functional Analysis*. McGraw-Hill, Inc.
- Silverman, B.W. (1986) *Density estimation for statistics and data analysis*. Chapman and Hall.

TABLE 1
Means and Standard Deviations of WLADE for
model (4.1) with t(2) innovations

		n=200		n=400		
ϕ_1	θ_1	ϕ_1	θ_1	ϕ_1	θ_1	
0.4	0.7	β	0.4092	0.6982	0.4078	0.6959
		SD	0.1013	0.0831	0.0716	0.0537
		AD	0.1156	0.0846	0.0779	0.0574
0.3	0.5	β	0.3019	0.5038	0.3009	0.4970
		SD	0.1153	0.1101	0.0916	0.0842
		AD	0.1165	0.1144	0.0926	0.0848
-0.5	-0.5	β	-0.4928	-0.4952	-0.4992	-0.4959
		SD	0.1075	0.1044	0.0705	0.0799
		AD	0.1164	0.1201	0.0779	0.0807

TABLE 2
Means and Standard Deviations of WLADE for
model (4.1) with Cauchy innovations

		n=200		n=400		
ϕ_1	θ_1	ϕ_1	θ_1	ϕ_1	θ_1	
0.4	0.7	β	0.4087	0.6918	0.4095	0.6924
		SD	0.0787	0.0535	0.0550	0.0382
		AD	0.0748	0.0530	0.0498	0.0379
0.3	0.5	β	0.3095	0.4917	0.3014	0.4936
		SD	0.0850	0.0776	0.0405	0.0360
		AD	0.0806	0.0749	0.0441	0.0371
-0.5	-0.5	β	-0.4994	-0.4925	-0.5009	-0.4974
		SD	0.0774	0.0849	0.0440	0.0498
		AD	0.0754	0.0831	0.0438	0.0478




Figure 1: Estimated values (\times) for the density functions of the normalized $\hat{\phi}_1$ (left panel) and $\hat{\theta}_1$ (right panel), together with their asymptotic limit — the $N(0,1)$ density (solid curves) for model (4.1) with t_2 innovations.




Figure 2: Estimated values (\times) for the density functions of the normalized $\hat{\phi}_1$ (left panel) and $\hat{\theta}_1$ (right panel), together with their asymptotic limit — the $N(0,1)$ density (solid curves) for model (4.1) with Cauchy innovations.




Figure 3: Boxplots of the AAE of LADE, WLADE, Whittle estimator, QMLE and Gauss-Newton estimator for model (4.1).




Figure 4: Boxplots of the AAE of WLADE and Gauss-Newton estimator for model (4.1) for large sample case.




Figure 5: Boxplots of AAE of WLADe with different α , γ and d for model (4.1) .




Figure 6: Simulation results for the Wald test. Differences between approximated P -value and the nominal significance level is plotted against the nominal level. The data were generated from (4.1) with $(\phi_1, \theta_1) = (0.3, 0)$ for the three panels on the left, and $(\phi_1, \theta_1) = (0.3, 0.5)$ for the three panels on the right.